

## **Patterns In Working Days Lost By Parents Of Children Newly-Diagnosed With Type 1 Diabetes (T1D)**

Ridyard, Colin; Blair, Jo ; Hughes, Dyfrig

### **Value in Health**

DOI:

[10.1016/j.jval.2017.08.1086](https://doi.org/10.1016/j.jval.2017.08.1086)

Published: 20/10/2017

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Ridyard, C., Blair, J., & Hughes, D. (2017). Patterns In Working Days Lost By Parents Of Children Newly-Diagnosed With Type 1 Diabetes (T1D). *Value in Health*, 20(9), A590. <https://doi.org/10.1016/j.jval.2017.08.1086>

#### **Hawliau Cyffredinol / General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## PATTERNS IN WORKING DAYS LOST BY PARENTS OF CHILDREN NEWLY-DIAGNOSED WITH TYPE 1 DIABETES (T1D).

**Ridyard CH**<sup>1</sup>, Blair J<sup>2</sup>, Hughes DA<sup>1</sup>, on behalf of the SCIPi Trial Investigators .<sup>3</sup>

<sup>1</sup>Bangor University, Bangor, United Kingdom, <sup>2</sup>Alder Hey Children's NHS Foundation Trust, Liverpool, United Kingdom, <sup>3</sup>Clinical Trials Research Centre, Liverpool, United Kingdom

**OBJECTIVES:** Continuous subcutaneous insulin infusions (CSII) are an alternative to multiple daily injections (MDI) for glycaemic control and reducing the risk of developing long term microvascular and macrovascular complications in type 1 diabetes (T1D). The objective of this study as part of a randomised clinical trial (SCIPi, ISRCTN29255275) was to assess patterns of work-related absences, and whether a difference was apparent between treatment groups. **METHODS:** Patients between 7 months and 15 years of age, newly diagnosed with T1D were eligible to participate in this pragmatic, open, multicentre, parallel group, randomised, controlled trial. Parental work-related absences in days from the preceding 3-months were measured by parental interview at randomisation, 3, 6, 9 and 12 month intervals. **RESULTS:** Actual time taken off work was reported by parents and/or guardians for 78% of participants (CSII=113, MDI=117). Absence from work between randomisation and 12-month follow-up in parents of patients in the CSII group was 3.6 days (95% CI 2.3, 4.9) compared to 2.7 days (1.9 to 3.4) in the MDI group [difference in means of 0.9 days (95% CI -0.6, 2.5)]. In the 3-month period prior to randomisation, however, absence from work in parents of patients in the CSII group was 5.5 days (95% CI 4.7, 6.3) compared to 4.9 days (4.1 to 5.7) in the MDI group [difference in means of 0.6 days (95% CI -0.5, 1.7)]. **CONCLUSIONS:** T1D diagnosis for a child can have a temporary disruptive influence on parental work commitments. However, there is no evidence to suggest that one treatment is associated with any more or less work-related absences than the other.